



EUROPEAN CENTRAL BANK

WORKING PAPER SERIES

NO. 487 / MAY 2005

**COMPUTING
SECOND-ORDER-ACCURATE
SOLUTIONS FOR RATIONAL
EXPECTATION MODELS
USING LINEAR SOLUTION
METHODS**

by Giovanni Lombardo
and Alan Sutherland

ECB EZB EKT EKP





EUROPEAN CENTRAL BANK



WORKING PAPER SERIES

NO. 487 / MAY 2005

COMPUTING SECOND-ORDER-ACCURATE SOLUTIONS FOR RATIONAL EXPECTATION MODELS USING LINEAR SOLUTION METHODS ¹

by Giovanni Lombardo ²
and Alan Sutherland ³

In 2005 all ECB publications will feature a motif taken from the €50 banknote.

This paper can be downloaded without charge from <http://www.ecb.int> or from the Social Science Research Network electronic library at http://ssrn.com/abstract_id=711167.

¹ In writing this paper we benefited from comments from Thomas Werner, Pierpaolo Benigno, Massimo Rostagno and Steve Ambler. Clearly, the content of this paper does not necessarily reflect their views. The content of this paper does not necessarily reflect the views of the European Central Bank.

² European Central Bank, Kaiserstrasse 29, 60311 Frankfurt am Main, Germany; e-mail: giovanni.lombardo@ecb.int

³ School of Economics and Finance, University of St Andrews, St Andrews, KY16 9AL, United Kingdom; e-mail: asj10@st-and.ac.uk

© European Central Bank, 2005

Address

Kaiserstrasse 29
60311 Frankfurt am Main, Germany

Postal address

Postfach 16 03 19
60066 Frankfurt am Main, Germany

Telephone

+49 69 1344 0

Internet

<http://www.ecb.int>

Fax

+49 69 1344 6000

Telex

411 144 ecb d

All rights reserved.

Reproduction for educational and non-commercial purposes is permitted provided that the source is acknowledged.

The views expressed in this paper do not necessarily reflect those of the European Central Bank.

The statement of purpose for the ECB Working Paper Series is available from the ECB website, <http://www.ecb.int>.

ISSN 1561-0810 (print)
ISSN 1725-2806 (online)

CONTENTS

Abstract	4
Non-technical summary	5
1 Introduction	6
2 A two-step solution method	7
3 State-space solutions to steps 1 and 2	9
3.1 Step 1	9
3.2 Step 2	11
4 An example: the neoclassical growth model	12
5 Conclusion	15
Appendix	15
References	21
European Central Bank working paper series	23

Abstract

This paper shows how to compute a second-order accurate solution of a non-linear rational expectation model using algorithms developed for the solution of *linear* rational expectation models. The result is a state-space representation for the realized values of the variables of the model. This state-space representation can easily be used to compute impulse responses as well as conditional and unconditional forecasts.

JEL classification: C63, E0.

Keywords Second order approximation; Solution method for rational expectation models.

Non-technical summary

Recently, the traditional application of the linearisation approach to the solution of dynamic general equilibrium models has shown some important limitations. Uncertainty and rational expectations are two of the most characteristic assumptions adopted in modern macroeconomic models. These assumptions have implications regarding the dynamics of the model economy as well as regarding the average level around which the economic variables are expected to fluctuate. The linearisation approach, by approximating the non-linear structural model by linear equations, is not able to take fully into account the role of uncertainty. The linearisation approach imposes certainty equivalence on a model so that some of the stochastic properties of the non-linear model are lost. On the contrary, a non-linear (e.g. quadratic) approximation of the model does not impose certainty equivalence on the economic relationships and provides a better measure of the effects of uncertainty on the economic variables. Furthermore, at least in some cases, approximations of order larger than one could also improve the accuracy of the solution.

In particular, recent developments in the analysis of monetary and fiscal policy have shown that a better characterization of the policy problem can be obtained by taking (at least) a second order expansion of the model around some point of interest (e.g. the non-stochastic steady state of the model).

One major advantage of the linearisation approach is that it requires only the use of linear algebra which makes it readily implementable on computers. This paper shows that also a second-order expansion of the non-linear model can be solved by using only linear algebra rules that are widely adopted in economics and econometrics textbooks. We show, in particular, that the same solution algorithms (and computer codes) that have been extensively used to solve linear-rational-expectation models can be used to solve second-order expansions of non-linear models. The result is the familiar state-space representation that is commonly associated with linear-rational-expectation models.

1 Introduction

This paper shows how algorithms devised for the solution of linear rational expectation models can be effectively employed to solve non-linear rational expectation models that are approximated to the second order of accuracy. Currently, researchers can choose from a number of algorithms for the solution of linear rational expectation models, i.e. models approximated to the first order of accuracy. An incomplete list would include direct methods like Blanchard and Kahn (1980), Sims (2000a) and Klein (2000) and methods based on the undetermined coefficients technique like Uhlig (1999) and Christiano (1998). At the same time a growing macroeconomic literature is addressing issues that can be studied only by taking into account (at least) the second-order terms of the rational expectation models. The welfare-based monetary policy analysis in Woodford (2003) is emblematic of this new focus. A number of papers describe how to derive the second-order expansion of rational expectation models and how to solve the approximated system. A non-exhaustive list should include Schmitt-Grohé and Uribe (2004), Jin and Judd (2002), Sims (2000b), Kim and Kim (2003), Kim *et al* (2003), Benigno and Woodford (2004a, 2004b) and Sutherland (2002). Most of these papers are associated with computer algorithms devised to solve the second-order-approximated models.¹ Yet, these algorithms (with the exception of Sutherland (2002)) are different from those used to solve linear rational expectation models. Furthermore, their description is often very heavy in terms of notation (e.g. they make use of the “tensor” notation).

In this paper we show that second-order accurate state-space solutions can be obtained simply by use of algorithms devised for linear rational expectations models. An important aspect of the method we propose is that it can be described using standard linear algebra notation, of the same type that would be used in linear rational expectations models (as described, for instance, in Ljungqvist and Sargent (2000)).² The basic structure of the solution technique employed in this paper follows the method suggested by Sutherland (2002). Nevertheless, our paper makes two important extensions to the results shown in Sutherland (2002). Firstly, we are able to derive

¹Benigno and Woodford (2004a, 2004b) represent an exception since their aim is to give an analytical solution to the model. Their approach is nevertheless very similar to that followed by Sutherland (2002). The general method proposed by Sutherland (2002) was developed independently but is similar to the procedure adopted by Canton (1996) in the context of a specific model.

²See Juillard (2003) for a “concise” formulation of the perturbation method that relies more heavily on matrix algebra.

second-order accurate solutions in *state-space form*. Secondly, we derive second-order accurate solutions for the *realized values* of the variables (as opposed to their conditional forecast). Thus, contrary to what is stated in Sutherland (2002), the two-step solution method described here is as general as any other second-order accurate solution method currently available in the literature (including those described by Schmitt-Grohé and Uribe (2004) and Sims (2000b)).

This paper is organized as follows. In Section 2 we outline the basic structure of the two-step solution procedure. In Section 3 the state-space form of the solutions to each step are described in more detail. Section 4 applies the solution method to the simple neoclassical growth model. This is a convenient benchmark which is used by both Sutherland (2002) and Schmitt-Grohé and Uribe (2004). Section 5 concludes.

2 A Two-Step Solution Method

It is assumed that the second-order approximation of the equations of a model can be written in the following matrix form³

$$A_1 \begin{bmatrix} s_{t+1} \\ E_t[c_{t+1}] \end{bmatrix} = A_2 \begin{bmatrix} s_t \\ c_t \end{bmatrix} + A_3 x_t + A_4 \Lambda_t + A_5 E_t[\Lambda_{t+1}] + O(\epsilon^3) \quad (1)$$

$$x_t = N x_{t-1} + \varepsilon_t \quad (2)$$

$$\Lambda_t = \text{vech} \left(\begin{bmatrix} x_t \\ s_t \\ c_t \end{bmatrix} \begin{bmatrix} x_t & s_t & c_t \end{bmatrix} \right) \quad (3)$$

where s is a vector of predetermined variables (i.e. $E_t[s_{t+1}] = s_{t+1}$), c is a vector of jump variables, x is a vector of exogenous forcing processes, ε is a vector of i.i.d. shocks. Λ_t is a vector of all the squares and cross-products

³The second-order approximation of a model is generated by replacing each side of each equation with a second-order Taylor series expansion around an appropriate point of approximation. It is usually convenient to approximate around a non-stochastic steady state. It is also usually convenient to measure variables as log-deviations from this non-stochastic steady state.

It is important to note that, in taking second-order approximations, expectations operators should be preserved in the positions they arise in the non-approximated model. This is because (unlike the case of first-order approximation) certainty equivalence can not be assumed in the second-order approximated model.

of the variables of the model.⁴ $A_1.. A_5$ are matrices of coefficients, E_t is the expectations operator conditional on information at time t and $O(\epsilon^3)$ contains all terms which are of order three or higher in deviations from the point of approximation.⁵

The objective is to use (1) to derive second-order accurate time paths of s and c . The solution method described in this paper is based on the following two observations: (i) second-order accurate solutions to (1) can be obtained using purely linear methods if a second-order accurate solution for the time-path of Λ is known; and (ii) a second-order accurate solution for the time path of Λ can itself be obtained using purely linear solution methods.

The first observation is self-evidently true. If the time path of Λ is known then (1) can be regarded as a linear rational expectations system with exogenous forcing processes Λ and x . Such a system can be solved using any standard linear solution method.

The second observation is less obvious. To understand (ii) notice that terms of order two and above in the behaviour of x , s and c become terms of order three and above in the squares and cross products of x , s and c . It must therefore follow that the second-order accurate behaviour of Λ depends only on the first-order accurate behaviour of x , s and c . Thus it is possible to generate second-order accurate solutions for Λ by considering first-order accurate solutions for x , s and c . First-order accurate solutions for x , s and c can easily be obtained by solving the linear system

$$A_1 \begin{bmatrix} s_{t+1} \\ E_t [c_{t+1}] \end{bmatrix} = A_2 \begin{bmatrix} s_t \\ c_t \end{bmatrix} + A_3 x_t + O(\epsilon^2) \quad (4)$$

which is derived from the first-order terms in (1). Here $O(\epsilon^2)$ contains all terms of order two and above in deviations from the non-stochastic steady state of the model.

It is now simple to state the two-step solution process.

Step 1: *Use the first-order dynamic system (4) to derive a second-order accurate solution for Λ .*

⁴The cross-products could involve variables with different time subscripts. By using the state-space solution discussed below, these cross-products can be easily reduced to products between contemporaneous realizations of the variables, i.e. Λ_t . See the Appendix for an explanation of the *vech* notation.

⁵It is assumed the distribution and dynamics of the exogenous driving processes in the model are such that no x variable can ever deviate from its deterministic steady state by more than ϵ .

Step 2: Use the solution for Λ derived in step 1 and the second-order dynamic system (1) to derive second-order accurate solutions for s and c .

An important difference between the current paper and Sutherland (2002) is that in Step 1 we are able to derive a linear state-space representation of the realised behaviour of Λ . The combination of this linear state-space representation of the dynamics of Λ and (1) yields an augmented system where the dynamics of Λ are treated as an additional set of linear exogenous forcing processes. Thus the non-linear system (1) is recast as a purely linear system with linear forcing processes. The solution to Step 2 can therefore also be written in a simple state-space form which can be used to generate second-order accurate impulse responses or second-order accurate values for conditional first and second moments at any horizon.

3 State-Space Solutions to Steps 1 and 2

In this section we describe the state-space solutions to Steps 1 and 2 in more detail and show explicitly how the second-order (i.e. non-linear) problem can be solved using purely linear solution methods. In this section we stress that what matters is the state-space representation of the solutions, not the particular algorithm used to derive the solutions. In the Appendix we describe in more detail how the QZ decomposition (as described in Klein (2000)) can be used to derive state-space solutions to each step. Matlab codes which implement the solution algorithm described in the Appendix are available from the authors.

3.1 Step 1

The first-order representation of our system (4) can be solved using any standard linear rational expectations method to yield a state-space representation of the following form

$$s_t^f = F_1 x_{t-1} + F_2 s_{t-1}^f \quad (5)$$

$$c_t^f = P_1 x_t + P_2 s_t^f \quad (6)$$

where the superscript ‘ f ’ indicates that these are first-order accurate solutions.⁶ It is convenient to rewrite this solution in a more compact form as

$$\begin{bmatrix} x_t^f \\ s_t^f \\ c_t^f \end{bmatrix} = \Omega \begin{bmatrix} x_t \\ s_t^f \end{bmatrix} \quad (7)$$

$$\begin{bmatrix} x_t \\ s_t^f \end{bmatrix} = \Phi \begin{bmatrix} x_{t-1} \\ s_{t-1}^f \end{bmatrix} + \Gamma \varepsilon_t \quad (8)$$

where

$$\Omega = \begin{bmatrix} I & 0 \\ 0 & I \\ P_1 & P_2 \end{bmatrix}, \quad \Phi = \begin{bmatrix} N & 0 \\ F_1 & F_2 \end{bmatrix}, \quad \Gamma = \begin{bmatrix} I \\ 0 \end{bmatrix} \quad (9)$$

Using the matrices L^c and L^h such that⁷

$$\begin{aligned} \text{vech}(\cdot) &= L^c \text{vec}(\cdot) \\ L^h \text{vech}(\cdot) &= \text{vec}(\cdot) \end{aligned}$$

it is easy to see that

$$\Lambda_t = R V_t \quad (10)$$

$$V_t = \tilde{\Phi} V_{t-1} + \tilde{\Gamma} \tilde{\varepsilon}_t + \tilde{\Psi} \tilde{\xi}_t \quad (11)$$

where $R = L^c (\Omega \otimes \Omega) L^h$, $\tilde{\Phi} = L^c (\Phi \otimes \Phi) L^h$, $\tilde{\Gamma} = L^c (\Gamma \otimes \Gamma) L^h$, and $\tilde{\varepsilon}_t = \text{vech}(\varepsilon_t \varepsilon_t')$, and where

$$V_t = \text{vech} \left(\begin{bmatrix} x_t \\ s_t^f \end{bmatrix} \begin{bmatrix} x_t & s_t^f \end{bmatrix} \right),$$

$$\tilde{\Psi} = L^c [(\Phi \otimes \Gamma) + (\Gamma \otimes \Phi) P']$$

$$\tilde{\xi}_t = \text{vec} \left(\begin{bmatrix} x_{t-1} \\ s_{t-1}^f \end{bmatrix} \varepsilon_t' \right)$$

(See the Appendix for a definition of the \otimes operator and also a discussion of the derivation of the ‘permutation’ matrix P .) Thus a second-order accurate representation of the dynamics of Λ can be written as a self-contained system in state-space form.

⁶Henceforth to simplify notation the term representing the approximation residual is omitted from all equations.

⁷Note that $L^h L^c = I$. See Hamilton (1996, p 300-302). Note also that the use of these matrices is not necessary in order to solve the model. Indeed one could simply vectorize the variance covariance dynamic system (use vec instead of vech). The suggested representation is clearly dictated by efficiency reasons.

3.2 Step 2

We can now use equation (10) to substitute out Λ_t and Λ_{t+1} in equation (1). This gives a new augmented form for the second-order accurate representation of the model as follows

$$A_1 \begin{bmatrix} s_{t+1} \\ E_t[c_{t+1}] \end{bmatrix} = A_2 \begin{bmatrix} s_t \\ c_t \end{bmatrix} + A_3 x_t + G V_t + H \Sigma \quad (12)$$

$$V_t = \tilde{\Phi} V_{t-1} + \tilde{\Gamma} \tilde{\varepsilon}_t + \tilde{\Psi} \tilde{\xi}_t \quad (13)$$

$$x_t = N x_{t-1} + \varepsilon_t \quad (14)$$

$$s_t^f = F_1 x_{t-1} + F_2 s_{t-1}^f \quad (15)$$

where⁸

$$G = (A_4 R + A_5 R \tilde{\Phi}), \quad H = A_5 R \tilde{\Gamma}, \quad \Sigma = E_t \tilde{\varepsilon}_{t+1} \quad (16)$$

The important point to notice is that this new representation of the second-order approximation of the model can now be solved in state-space form using any linear rational expectations solution method.⁹ A state-space representation of the solution to this system is the following

$$s_t = F_1 x_{t-1} + F_2 s_{t-1} + F_3 V_{t-1} + F_4 \Sigma \quad (17)$$

$$c_t = P_1 x_t + P_2 s_t + P_3 V_t + P_4 \Sigma \quad (18)$$

$$V_t = \tilde{\Phi} V_{t-1} + \tilde{\Gamma} \tilde{\varepsilon}_t + \tilde{\Psi} \tilde{\xi}_t \quad (19)$$

$$x_t = N x_{t-1} + \varepsilon_t \quad (20)$$

$$s_t^f = F_1 x_{t-1} + F_2 s_{t-1}^f \quad (21)$$

For any given initial conditions for s , V and x , this state-space system can be used to generate second-order accurate impulse responses to the exogenous shocks.¹⁰ It can also be used to generate second-order accurate stochastic simulations for computer generated random realisations of the innovations.

Furthermore, the state-space representation provides a convenient way to calculate second-order accurate solutions for conditional first and second

⁸Note that $E_t[\tilde{\xi}_{t+1}] = 0$.

⁹This is despite the presence of the cross-product term $\tilde{\xi}_t$. The cross-product term is zero in expectation and therefore does not affect the forward-looking dynamics of the model. The forward-looking dynamics of the model are therefore entirely linear.

¹⁰Notice that, in this case, the cross product term $\tilde{\xi}_t$ is zero in all periods because x_{t-1} and s_{t-1}^f are zero in the first period of the impulse response simulation and ε_t is zero in all periods other than the first period of the impulse response simulation. Equation (21) is therefore not relevant for generating an impulse response solution.



moments for the time-paths for the variables of the model. By simply applying the conditional expectation operator through all the equations in (17) to (21) we can compute first and second conditional moments at all horizons.¹¹

4 An Example: The Neoclassical Growth Model

As an example of the use of the above algorithm consider the simple neoclassical growth model consisting of three equations: an Euler consumption (c) equation, a capital (k) accumulation equation and an i.i.d. process for the (log) of the productivity shock (a).¹² That is

$$c_t^{-\gamma} = \alpha\beta E_t [a_{t+1}k_{t+1}^{\alpha-1}c_{t+1}^{-\gamma}] \quad (22)$$

$$k_{t+1} = a_t k_t^\alpha - c_t \quad (23)$$

$$\hat{a}_t \equiv \log a_t = \varepsilon_t \quad (24)$$

The equation-by-equation second-order Taylor expansion of this simple model is as follows (where hats indicate log-deviations from a non-stochastic steady state).

$$\begin{aligned} -\gamma\hat{c}_t + (1/2)\gamma^2\hat{c}_t^2 &= -\gamma E_t \hat{c}_{t+1} + (\alpha - 1)\hat{k}_{t+1} + \\ &\quad (1/2)E_t \left[\left(\hat{a}_{t+1} + \gamma\hat{c}_{t+1} + (\alpha - 1)\hat{k}_{t+1} \right)^2 \right] \end{aligned} \quad (25)$$

$$\begin{aligned} \theta\hat{k}_{t+1} + (1/2)\theta\hat{k}_{t+1}^2 &= \hat{a}_t + \alpha\hat{k}_t - \phi\hat{c}_t - (1/2)\phi\hat{c}_t^2 + \\ &\quad (1/2)\alpha^2\hat{k}_t^2 + (1/2)\hat{a}_t^2 + \alpha\hat{a}_t\hat{k}_t \end{aligned} \quad (26)$$

$$\hat{a}_t = \varepsilon_t \quad (27)$$

¹¹An increasing number of macroeconomic papers make use of second-order approximation methods in order to analyze the welfare effects of fiscal and monetary policies as well as in order to derive optimal policies. This requires solutions for first and second moments rather than solutions for realised values. This is in fact the main focus of Sutherland (2002) and Benigno and Woodford (2004a, 2004b). Notice that the cross-product term, ξ_t , is irrelevant for generating expected paths because it is zero in expectation. Equation (21) is therefore also irrelevant in this case.

¹²This model corresponds to one of the examples used by Schmitt-Grohé and Uribe (2004). The assumption of zero persistence in the productivity shock and no depreciation in the capital stock are also made in Schmitt-Grohé and Uribe (2004). These assumptions are made for simplicity only and are not required for the application of the solution algorithm.

where $\phi = \frac{c_{ss}}{c_{ss}+k_{ss}}$, $\theta = \frac{k_{ss}}{c_{ss}+k_{ss}}$. The approximation-error term is not shown for simplicity.¹³ Equations (22), (23) and (24) are obtained by replacing each side of equations (25), (26) and (27) with a second-order (logarithmic) Taylor series expansion around the non-stochastic steady state. Notice that the conditional expectations operator which appears in (22) is preserved in equation (25).¹⁴

Next, we cast the model in matrix notation as follows

$$A_1 \begin{bmatrix} \hat{k}_{t+1} \\ E_t[\hat{c}_{t+1}] \end{bmatrix} = A_2 \begin{bmatrix} \hat{k}_t \\ \hat{c}_t \end{bmatrix} + A_3 a_t + A_4 \Lambda_t + A_5 E_t[\Lambda_{t+1}] \quad (28)$$

where

$$\Lambda_t = \begin{bmatrix} \hat{a}_t^2 & \hat{a}_t \hat{k}_t & \hat{k}_t^2 & \hat{a}_t \hat{c}_t & \hat{k}_t \hat{c}_t & \hat{c}_t^2 \end{bmatrix}$$

$$A_1 = \begin{bmatrix} \theta & 0 \\ 1 - \alpha & \gamma \end{bmatrix} \quad A_2 = \begin{bmatrix} \alpha & -\phi \\ 0 & \gamma \end{bmatrix} \quad A_3 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$A_4 = \begin{bmatrix} 1/2 & \alpha & \alpha^2/2 & 0 & 0 & -\phi/2 \\ 0 & 0 & 0 & 0 & 0 & -\gamma^2/2 \end{bmatrix}$$

$$A_5 = \begin{bmatrix} 0 & 0 & \theta/2 & 0 & 0 & 0 \\ 1/2 & \alpha - 1 & (\alpha - 1)^2/2 & \gamma & \gamma(\alpha - 1) & \gamma^2/2 \end{bmatrix}$$

The following parameter values are used: $\gamma = 2$, $\alpha = 0.3$, $\beta = 0.95$, $\theta = 0.285$, $\phi = 0.715$.

We are now ready to use the two-step algorithm outlined above. Step 1 of the algorithm yields the following state-space representation for the evolution

¹³Nevertheless, it is useful to recall that this is a *local* approximation and hence the error term might be large for large departures from the approximation point (the steady state in our case) (see Jin and Judd (2002) for a discussion of the importance of the *local* perspective in this kind of exercises).

¹⁴Note that, by definition, $E_t[k_{t+1}] = k_{t+1}$ and $E_t[a_{t+1}] = 0$.

of Λ_t (i.e. equations (10) and (11)):¹⁵

$$\begin{bmatrix} \hat{a}_t^2 \\ \hat{a}_t \hat{k}_t^f \\ (\hat{k}_t^f)^2 \\ \hat{a}_t \hat{c}_t^f \\ \hat{k}_t^f \hat{c}_t^f \\ (\hat{c}_t^f)^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0.84174 & 0.25252 & 0 \\ 0 & 0.84174 & 0.25252 \\ 0.70853 & 0.42512 & 0.063768 \end{bmatrix} \begin{bmatrix} \hat{a}_t^2 \\ \hat{a}_t \hat{k}_t^f \\ (\hat{k}_t^f)^2 \end{bmatrix} \quad (29)$$

$$\begin{bmatrix} \hat{a}_t^2 \\ \hat{a}_t \hat{k}_t^f \\ (\hat{k}_t^f)^2 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1.9517 & 1.171 & 0.17565 \end{bmatrix} \begin{bmatrix} \hat{a}_{t-1}^2 \\ \hat{a}_{t-1} \hat{k}_{t-1}^f \\ (\hat{k}_{t-1}^f)^2 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} [\varepsilon_t^2] \\ + \begin{bmatrix} 0 & 0 \\ 1.397 & 0.41911 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \hat{a}_{t-1} \\ \hat{k}_{t-1}^f \end{bmatrix} \varepsilon_t \quad (30)$$

$$\begin{bmatrix} \hat{a}_t \\ \hat{k}_t^f \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1.397 & 0.41911 \end{bmatrix} \begin{bmatrix} \hat{a}_{t-1} \\ \hat{k}_{t-1}^f \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \end{bmatrix} [\varepsilon_t] \quad (31)$$

Step 2 of the algorithm yields the following state-space representation of the second-order accurate solution of the model:

$$\begin{bmatrix} \hat{k}_{t+1} \\ \hat{c}_t \end{bmatrix} = \begin{bmatrix} 1.397 & 0.41911 \\ 0.84174 & 0.25252 \end{bmatrix} \begin{bmatrix} \hat{a}_t \\ \hat{k}_t \end{bmatrix} \\ + \frac{1}{2} \begin{bmatrix} -0.077802 & -0.046681 & -0.0070022 \\ -0.056866 & -0.034120 & -0.005118 \end{bmatrix} \begin{bmatrix} \hat{a}_t^2 \\ \hat{a}_t \hat{k}_t^f \\ (\hat{k}_t^f)^2 \end{bmatrix} \quad (32) \\ + \frac{1}{2} \begin{bmatrix} 0.4820 \\ -0.1921 \end{bmatrix} \sigma^2$$

These numbers are identical to those reported in Schmitt-Grohé and Uribe (2004) for the same model.

Schmitt-Grohé and Uribe (2004) report results relating to two other models. We have applied our algorithm to both these other examples and have confirmed that it generates identical results to those reported by Schmitt-Grohé and Uribe (2004).

¹⁵In what follows \hat{k}^f and \hat{c}^f denote first-order accurate solutions for capital and consumption while \hat{k} and \hat{c} denote second-order accurate solutions for capital and consumption.

5 Conclusion

In this paper we have shown how a non-linear rational expectation model, approximated to the second order of accuracy, can be recast as a linear structure which can be solved in state-space form by means of standard algorithms developed for the solution of linear rational expectation models. This state-space form can be used to produce second-order accurate impulse responses as well as conditional and unconditional forecasts. We suggest that our algorithm is a convenient alternative to other second-order accurate solution methods proposed in recent literature. Compared to other methods, our algorithm seem to require a much more modest departure from the existing techniques used in dynamic-rational-expectations macroeconomics.

Appendix

Glossary of Matrix Algebra Notation and Rules

1. $\text{vec}(X)$: Vectorization. All columns of the $m \times n$ matrix X are stacked one under the other (starting with the first column).
2. $\text{vech}(X)$: As above except that only the upper triangular part of X is considered. Note that it is possible to construct a matrix L such that $L \text{vech} = \text{vec}$. Then, $(L' L)^{-1} L' \text{vec}(X) = \text{vech}(X)$.
3. \otimes : Kronecker product. E.g. $Z = X \otimes Y$. The elements of Z are the product of each element of X with the matrix Y .
4. **Vectorization of a product of matrices:** $\text{vec}(X Y Z) = (Z' \otimes X) \text{vec}(Y)$
5. **The permutation matrix P** Here we show how to construct the permutation matrix P such that $\text{vec}(Z) = P \text{vec}(Z')$. We start by noticing that the element $z_{i,j}$ of the generic matrix Z of dimension $m \times n$ will coincide with the element $z_{i+m(j-1)}^v$ in the vector $z^v = \text{vec}(Z)$, while it will coincide with the element $z_{j+n(i-1)}^v$ in the vector $\bar{z}^v = \text{vec}(Z')$. This information can be used to generate the matrix P . Generate an $m \times n$ matrix S such that $S = \text{vec}^{-1}([1, 2 \dots (m \cdot n)]')$, and an identity matrix I of dimension $m n \times m n$. Finally, the permutation matrix P is given by $P = I(:, \text{vec}(S'))$.

State-Space Solution to the First-Order System

Consider the first-order system

$$A_1 E_t \begin{bmatrix} s_{t+1} \\ c_{t+1} \end{bmatrix} = A_2 \begin{bmatrix} s_t \\ c_t \end{bmatrix} + A_3 x_t \quad (33)$$

$$x_t = N x_{t-1} + \varepsilon_t \quad (34)$$

By applying the QZ decomposition (Generalized Schur Decomposition) we can factorize the matrices A_1 and A_2 into

$$qA_1 z = \begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix}, \quad qA_2 z = \begin{bmatrix} b_{11} & b_{12} \\ 0 & b_{22} \end{bmatrix}$$

where matrix z has the property $z z' = I$. Hence

$$\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} E_t \begin{bmatrix} y_{1,t+1} \\ y_{2,t+1} \end{bmatrix} = \begin{bmatrix} b_{11} & b_{12} \\ 0 & b_{22} \end{bmatrix} \begin{bmatrix} y_{1,t} \\ y_{2,t} \end{bmatrix} + \begin{bmatrix} C_1 \\ C_2 \end{bmatrix} x_t \quad (35)$$

where

$$\begin{bmatrix} y_{1,t} \\ y_{2,t} \end{bmatrix} = \begin{bmatrix} z'_{11} & z'_{21} \\ z'_{12} & z'_{22} \end{bmatrix} \begin{bmatrix} s_t \\ c_t \end{bmatrix}$$

and

$$\begin{bmatrix} C_1 \\ C_2 \end{bmatrix} = qA_3$$

Without loss of generality we can assume that the system (35) has been ordered so that $b_{22}^{-1} a_{22}$ has roots inside the unit circle. Then the lower part of system (35) can be isolated and solved forward to get (absent bubbles)

$$\hat{y}_{2,t} = - [b_{22}^{-1} C_2 + T b_{22}^{-1} C_2 N + T^2 b_{22}^{-1} C_2 N^2 + \dots] x_t \quad (36)$$

where

$$T = b_{22}^{-1} a_{22}$$

As long as the series converges we can solve for the endogenous variables as

$$y_{2,t} = -M x_t$$

where

$$vec(M) = [I - (N' \otimes T)]^{-1} vec(b_{22}^{-1} C_2)$$

See the Glossary at the start of this Appendix for a general statement of the rule used to derive this expression.¹⁶

Finally we have

$$\hat{y}_{2,t} \equiv z'_{12}s_t + z'_{22}c_t = -Mx_t$$

so that

$$c_t = P_1x_t + P_2s_t \quad (37)$$

where

$$P_1 = -z'^{-1}_{22}M, \quad P_2 = -z'^{-1}_{22}z'_{12}$$

As for the state variables, solving for the upper part of (35) yields

$$\underbrace{(a_{11}z'_{21} + a_{12}z'_{22}) P_1}_{R_1} E_t x_{t+1} + \underbrace{[(a_{11}z'_{11} + a_{12}z'_{12}) + (a_{11}z'_{21} + a_{12}z'_{22}) P_2]}_{R_2} E_t s_{t+1} = \underbrace{[(b_{11}z'_{21} + b_{12}z'_{22}) P_1 + C_1]}_{D_1} x_t + \underbrace{[(b_{11}z'_{11} + b_{12}z'_{12}) + (b_{11}z'_{21} + b_{12}z'_{22}) P_2]}_{D_2} s_t$$

Thus

$$E_t [R_1 x_{t+1} + R_2 s_{t+1}] = D_1 x_t + D_2 s_t$$

or

$$s_{t+1} = \underbrace{(R_2^{-1}D_1 - R_2^{-1}R_1N)}_{F_1} x_t + \underbrace{R_2^{-1}D_2}_{F_2} s_t$$

where we have made use of the fact that $E_t s_{t+1} = s_{t+1}$ (because s is a vector of predetermined variables).

To sum up, the solution to the dynamic system (33) is

$$s_t = F_1 x_{t-1} + F_2 s_{t-1} \quad (38)$$

$$c_t = P_1 x_t + P_2 s_t \quad (39)$$

$$x_t = N x_{t-1} + \varepsilon_t \quad (40)$$

This is the solution given in (5) and (6) in the main text.

¹⁶Klein (referring to King and Watson (2002)) describes a computationally more efficient method to compute M .

State-Space Solution to the Second-Order System

Consider now the augmented second-order system

$$A_1 \begin{bmatrix} s_{t+1} \\ E_t[c_{t+1}] \end{bmatrix} = A_2 \begin{bmatrix} s_t \\ c_t \end{bmatrix} + A_3 x_t + G V_t + H \Sigma \quad (41)$$

$$V_t = \tilde{\Phi} V_{t-1} + \tilde{\Gamma} \tilde{\varepsilon}_t + \tilde{\Psi} \tilde{\xi}_t \quad (42)$$

$$x_t = N x_{t-1} + \varepsilon_t \quad (43)$$

$$s_t^f = F_1 x_{t-1} + F_2 s_{t-1}^f \quad (44)$$

Define $\bar{V} = (I - \tilde{\Phi})^{-1} \tilde{\Gamma}$ then

$$E_t[V_{t+n}] = \bar{V} \Sigma + \tilde{\Phi}^n (V_t - \bar{V} \Sigma)$$

Applying the QZ decomposition yields

$$\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} E_t \begin{bmatrix} y_{1,t+1} \\ y_{2,t+1} \end{bmatrix} = \begin{bmatrix} b_{11} & b_{12} \\ 0 & b_{22} \end{bmatrix} \begin{bmatrix} \hat{y}_{1,t} \\ \hat{y}_{2,t} \end{bmatrix} + \begin{bmatrix} C_1 \\ C_2 \end{bmatrix} x_t \\ + \begin{bmatrix} \hat{G}_1 \\ \hat{G}_2 \end{bmatrix} V_t + \begin{bmatrix} \hat{H}_1 \\ \hat{H}_2 \end{bmatrix} \Sigma \quad (45)$$

where the matrices a , b , q and z are all identical to those defined in the previous section and

$$\begin{bmatrix} \hat{G}_1 \\ \hat{G}_2 \end{bmatrix} = qG, \quad \begin{bmatrix} \hat{H}_1 \\ \hat{H}_2 \end{bmatrix} = qH$$

Again the lower part of system (45) can be isolated and solved forward to yield

$$y_{2,t} = - \left[b_{22}^{-1} C_2 + T b_{22}^{-1} C_2 N + T^2 b_{22}^{-1} C_2 N^2 + \dots \right] x_t \\ - \left[b_{22}^{-1} \hat{G}_2 + T b_{22}^{-1} \hat{G}_2 \tilde{\Phi} + T^2 b_{22}^{-1} \hat{G}_2 \tilde{\Phi}^2 + \dots \right] (V_t - \bar{V} \Sigma) \\ - \left[I + T + T^2 + \dots \right] b_{22}^{-1} (\hat{G}_2 \bar{V} + \hat{H}_2) \Sigma \quad (46)$$

where

$$T = b_{22}^{-1} a_{22}$$

As long as the series converges we can solve for the endogenous variables as

$$y_{2,t} = -M_1 x_t - M_2 (V_t - \bar{V} \Sigma) - M_3 \Sigma$$

where

$$\begin{aligned} \text{vec}(M_1) &= [I - (N' \otimes T)]^{-1} \text{vec}(b_{22}^{-1}C_2) \\ \text{vec}(M_2) &= \left[I - \left(\tilde{\Phi}' \otimes T \right) \right]^{-1} \text{vec}(b_{22}^{-1}\hat{G}_2) \\ M_3 &= [I - T]^{-1} b_{22}^{-1}(\hat{G}_2\bar{V} + \hat{H}_2) \end{aligned}$$

Finally we have

$$y_{2,t} \equiv z'_{12}s_t + z'_{22}c_t = -M_1x_t - M_2(V_t - \bar{V}\Sigma) - M_3\Sigma$$

so that

$$c_t = P_1x_t + P_2s_t + P_3V_t + P_4\Sigma \quad (47)$$

where

$$\begin{aligned} P_1 &= -z'_{22}{}^{-1}M_1 \\ P_2 &= -z'_{22}{}^{-1}z'_{12} \\ P_3 &= -z'_{22}{}^{-1}M_2 \\ P_4 &= -z'_{22}{}^{-1}[M_3 - M_2\bar{V}] \end{aligned}$$

The solution for the state variables can be obtained by solving for the upper part of (45). This yields

$$\begin{aligned} &\underbrace{(a_{11}z'_{21} + a_{12}z'_{22})P_1}_{R_1} E_t x_{t+1} + \underbrace{[(a_{11}z'_{11} + a_{12}z'_{12}) + (a_{11}z'_{21} + a_{12}z'_{22})P_2]}_{R_2} E_t s_{t+1} \\ &+ \underbrace{(a_{11}z'_{21} + a_{12}z'_{22})P_3}_{R_3} E_t V_{t+1} + \underbrace{(a_{11}z'_{21} + a_{12}z'_{22})P_4}_{R_4} \Sigma = \\ &\underbrace{[(b_{11}z'_{21} + b_{12}z'_{22})P_1 + C_1]}_{D_1} x_t + \underbrace{[(b_{11}z'_{11} + b_{12}z'_{12}) + (b_{11}z'_{21} + b_{12}z'_{22})P_2]}_{D_2} s_t \\ &+ \underbrace{[(b_{11}z'_{21} + b_{12}z'_{22})P_3 + \hat{G}_1]}_{D_3} V_t + \underbrace{[(b_{11}z'_{21} + b_{12}z'_{22})P_4 + \hat{H}_1]}_{D_4} \Sigma \end{aligned}$$

Thus

$$R_1 N x_{t-1} + R_2 s_t + R_3 (\tilde{\Phi} V_{t-1} + \tilde{\Gamma} \Sigma) + R_4 \Sigma = D_1 x_{t-1} + D_2 s_{t-1} + D_3 V_{t-1} + D_4 \Sigma$$

or

$$\begin{aligned} s_t &= \underbrace{R_2^{-1}(D_1 - R_1 N)}_{F_1} x_{t-1} + \underbrace{R_2^{-1}D_2}_{F_2} s_{t-1} \\ &\quad + \underbrace{R_2^{-1}(D_3 - R_3)}_{F_3} V_{t-1} + \underbrace{R_2^{-1}(D_4 - R_4 - R_3 \tilde{\Gamma})}_{F_4} \Sigma \end{aligned}$$

To sum up, the solution to the second-order system (41) is

$$s_t = F_1 x_{t-1} + F_2 s_{t-1} + F_3 V_{t-1} + F_4 \Sigma \quad (48)$$

$$c_t = P_1 x_t + P_2 s_t + P_3 V_t + P_4 \Sigma \quad (49)$$

$$x_t = N x_{t-1} + \varepsilon_t \quad (50)$$

$$V_t = \tilde{\Phi} V_{t-1} + \tilde{\Gamma} \tilde{\varepsilon}_t + \tilde{\Psi} \tilde{\xi}_t \quad (51)$$

$$s_t^f = F_1 x_{t-1} + F_2 s_{t-1}^f \quad (52)$$

This is the state-space form of the second-order solution given in equations (17) to (22) in the main text.

Notice that the QZ decomposition only needs to be applied once in the two-step procedure. The matrices a, b, q and z are the same in both steps, as are the solutions for F_1, F_2, P_1 and P_2 .¹⁷

¹⁷Only in cases where the realised and expected dynamics differ would it be necessary to compute the QZ decomposition twice.

References

- [1] Benigno, P and M Woodford (2004a) “Optimal Monetary and Fiscal Policy: A Linear Quadratic Approach” in Gertler and Rogoff (eds) *NBER Macroeconomics Annual*, 18, 271-333.
- [2] Benigno, P and M Woodford (2004b) “Optimal Stabilisation Policy When Wages are Sticky: The Case of a Distorted Steady State” NBER Working Paper 10839.
- [3] Blanchard, O J and C M Kahn (1980) “The Solution to Linear Difference Models under Rational Expectations” *Econometrica*, 48, 1305-1311.
- [4] Canton, E (1996) “Business Cycles in a Two-Sector Model of Endogenous Growth” CentER Discussion Paper No 96116.
- [5] Christiano, L J (1998) “Solving Dynamic Equilibrium Models by a Method of Undetermined Coefficients” NBER Working Paper T0225.
- [6] Hamilton, J D (1994) *Time Series Analysis*, Princeton University Press, Princeton.
- [7] Jin, H-H and Judd, K J (2002) “Perturbation Methods for General Dynamic Stochastic Models” unpublished manuscript, Stanford University.
- [8] Juillard, M (2003) “Solving Stochastic Dynamic Models: A k-Order Perturbation Approach” unpublished manuscript, University Paris 8.
- [9] Kim, J, S Kim, E Schaumburg and C Sims (2003) “Calculating and Using Second Order Accurate Solutions of Discrete Time Dynamic Equilibrium Models” unpublished manuscript, Princeton University.
- [10] Kim, J and S H Kim (2003) ‘Spurious Welfare Reversals in International Business Cycle Models’, *Journal of International Economics*, 60, 471-500.
- [11] King, R G and M W Watson (2002) “System Reduction and Solution Algorithms for Singular Linear Difference Systems under Rational Expectations” *Computational Economics*, 20, 57-86.
- [12] Klein, P (2000) “Using the Generalised Schur Form to Solve a Multivariate Linear Rational Expectations Model” *Journal of Economic Dynamics and Control*, 24, 1405-1423.
- [13] Ljungqvist, L and T J. Sargent (2000) *Recursive Macroeconomic Theory*, MIT Press, Cambridge, MA.

- [14] Schmitt-Grohé, S and M Uribe (2004) “Solving Dynamic General Equilibrium Models using a Second-Order Approximation to the Policy Function”, *Journal of Economic Dynamics and Control*, 28, 755-75.
- [15] Sims, C (2000a) “Solving Linear Rational Expectations Models” unpublished manuscript, Princeton University.
- [16] Sims, C (2000b) “Second-Order Accurate Solutions of Discrete Time Dynamic Equilibrium Models” unpublished manuscript, Princeton University.
- [17] Sutherland, A (2002) “A Simple Second-Order Solution Method for Dynamic General Equilibrium Models” CEPR Discussion Paper No. 3554.
- [18] Uhlig, H (1999) “A Toolkit for Analysing Nonlinear Dynamic Stochastic Models Easily” in Marimon and Scott (eds) *Computational Methods for the Study of Dynamic Economies*, OUP, Oxford.
- [19] Woodford, M (2003) *Interest and Prices: Foundations of a Theory of Monetary Policy*, Princeton University Press, Princeton.

European Central Bank working paper series

For a complete list of Working Papers published by the ECB, please visit the ECB's website (<http://www.ecb.int>)

- 435 "Reforming public expenditure in industrialised countries: are there trade-offs?"
by L. Schuknecht and V. Tanzi, February 2005.
- 436 "Measuring market and inflation risk premia in France and in Germany"
by L. Cappiello and S. Guéné, February 2005.
- 437 "What drives international bank flows? Politics, institutions and other determinants"
by E. Papaioannou, February 2005.
- 438 "Quality of public finances and growth" by A. Afonso, W. Ebert, L. Schuknecht and M. Thöne,
February 2005.
- 439 "A look at intraday frictions in the euro area overnight deposit market"
by V. Brousseau and A. Manzanares, February 2005.
- 440 "Estimating and analysing currency options implied risk-neutral density functions for the largest
new EU member states" by O. Castrén, February 2005.
- 441 "The Phillips curve and long-term unemployment" by R. Llaudes, February 2005.
- 442 "Why do financial systems differ? History matters" by C. Monnet and E. Quintin, February 2005.
- 443 "Explaining cross-border large-value payment flows: evidence from TARGET and EURO I data"
by S. Rosati and S. Secola, February 2005.
- 444 "Keeping up with the Joneses, reference dependence, and equilibrium indeterminacy" by L. Stracca
and Ali al-Nowaihi, February 2005.
- 445 "Welfare implications of joining a common currency" by M. Ca' Zorzi, R. A. De Santis and F. Zampolli,
February 2005.
- 446 "Trade effects of the euro: evidence from sectoral data" by R. Baldwin, F. Skudelny and D. Taglioni,
February 2005.
- 447 "Foreign exchange option and returns based correlation forecasts: evaluation and two applications"
by O. Castrén and S. Mazzotta, February 2005.
- 448 "Price-setting behaviour in Belgium: what can be learned from an ad hoc survey?"
by L. Aucremanne and M. Druant, March 2005.
- 449 "Consumer price behaviour in Italy: evidence from micro CPI data" by G. Veronese, S. Fabiani,
A. Gattulli and R. Sabbatini, March 2005.
- 450 "Using mean reversion as a measure of persistence" by D. Dias and C. R. Marques, March 2005.
- 451 "Breaks in the mean of inflation: how they happen and what to do with them" by S. Corvoisier
and B. Mojon, March 2005.
- 452 "Stocks, bonds, money markets and exchange rates: measuring international financial transmission"
by M. Ehrmann, M. Fratzscher and R. Rigobon, March 2005.

- 453 “Does product market competition reduce inflation? Evidence from EU countries and sectors” by M. Przybyła and M. Roma, March 2005.
- 454 “European women: why do(n’t) they work?” by V. Genre, R. G. Salvador and A. Lamo, March 2005.
- 455 “Central bank transparency and private information in a dynamic macroeconomic model” by J. G. Pearlman, March 2005.
- 456 “The French block of the ESCB multi-country model” by F. Boissay and J.-P. Villetelle, March 2005.
- 457 “Transparency, disclosure and the Federal Reserve” by M. Ehrmann and M. Fratzscher, March 2005.
- 458 “Money demand and macroeconomic stability revisited” by A. Schabert and C. Stoltenberg, March 2005.
- 459 “Capital flows and the US ‘New Economy’: consumption smoothing and risk exposure” by M. Miller, O. Castrén and L. Zhang, March 2005.
- 460 “Part-time work in EU countries: labour market mobility, entry and exit” by H. Buddelmeyer, G. Mourre and M. Ward, March 2005.
- 461 “Do decreasing hazard functions for price changes make any sense?” by L. J. Álvarez, P. Burriel and I. Hernando, March 2005.
- 462 “Time-dependent versus state-dependent pricing: a panel data approach to the determinants of Belgian consumer price changes” by L. Aucremanne and E. Dhyne, March 2005.
- 463 “Break in the mean and persistence of inflation: a sectoral analysis of French CPI” by L. Bilke, March 2005.
- 464 “The price-setting behavior of Austrian firms: some survey evidence” by C. Kwapil, J. Baumgartner and J. Scharler, March 2005.
- 465 “Determinants and consequences of the unification of dual-class shares” by A. Pajuste, March 2005.
- 466 “Regulated and services’ prices and inflation persistence” by P. Lünemann and T. Y. Mathä, April 2005.
- 467 “Socio-economic development and fiscal policy: lessons from the cohesion countries for the new member states” by A. N. Mehrotra and T. A. Peltonen, April 2005.
- 468 “Endogeneities of optimum currency areas: what brings countries sharing a single currency closer together?” by P. De Grauwe and F. P. Mongelli, April 2005.
- 469 “Money and prices in models of bounded rationality in high inflation economies” by A. Marcet and J. P. Nicolini, April 2005.
- 470 “Structural filters for monetary analysis: the inflationary movements of money in the euro area” by A. Bruggeman, G. Camba-Méndez, B. Fischer and J. Sousa, April 2005.
- 471 “Real wages and local unemployment in the euro area” by A. Sanz de Galdeano and J. Turunen, April 2005.
- 472 “Yield curve prediction for the strategic investor” by C. Bernadell, J. Coche and K. Nyholm, April 2005.

- 473 “Fiscal consolidations in the Central and Eastern European countries” by A. Afonso, C. Nickel and P. Rother, April 2005.
- 474 “Calvo pricing and imperfect common knowledge: a forward looking model of rational inflation inertia” by K. P. Nimark, April 2005.
- 475 “Monetary policy analysis with potentially misspecified models” by M. Del Negro and F. Schorfheide, April 2005.
- 476 “Monetary policy with judgment: forecast targeting” by L. E. O. Svensson, April 2005.
- 477 “Parameter misspecification and robust monetary policy rules” by C. E. Walsh, April 2005.
- 478 “The conquest of U.S. inflation: learning and robustness to model uncertainty” by T. Cogley and T. J. Sargent, April 2005.
- 479 “The performance and robustness of interest-rate rules in models of the euro area” by R. Adalid, G. Coenen, P. McAdam and S. Siviero, April 2005.
- 480 “Insurance policies for monetary policy in the euro area” by K. Küster and V. Wieland, April 2005.
- 481 “Output and inflation responses to credit shocks: are there threshold effects in the euro area?” by A. Calza and J. Sousa, April 2005.
- 482 “Forecasting macroeconomic variables for the new member states of the European Union” by A. Banerjee, M. Marcellino and I. Masten, May 2005.
- 483 “Money supply and the implementation of interest rate targets” by A. Schabert, May 2005.
- 484 “Fiscal federalism and public inputs provision: vertical externalities matter” by D. Martínez-López, May 2005.
- 485 “Corporate investment and cash flow sensitivity: what drives the relationship?” by P. Mizen and P. Vermeulen, May 2005.
- 486 “What drives productivity growth in the new EU member states? The case of Poland” by M. Kolasa, May 2005.
- 487 “Computing second-order-accurate solutions for rational expectation models using linear solution methods” by G. Lombardo and A. Sutherland, May 2005.

